

Ripples in the Anterior Auditory Field and Inferior Colliculus of the Ferret

Didier Depireux Nina Kowalski

Shihab Shamma

Tony Owens Huib Versnel

Amitai Kohn

University of Maryland College Park


Supported in part by the Air Force Office of Scientific Research, the Office of Naval Research, the
National Science Foundation Grant CD-8803012 and the National Institutes of Health.

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 1996		2. REPORT TYPE		3. DATES COVERED 00-00-1996 to 00-00-1996	
4. TITLE AND SUBTITLE Ripples in the Anterior Auditory Field and Inferior Colliculus of the Ferret				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Office of Naval Research,One Liberty Center,875 North Randolph Street Suite 1425,Arlington,VA,22203-1995				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES The original document contains color images.					
14. ABSTRACT see report					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES 26	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			



Methods

- Responses of single units in AI, anterior field (AAF), and Inferior Colliculus (IC) in the barbiturate-anaesthetized ferret were recorded with single tungsten electrodes. Data were collected from a total of 7, 5 and 11 (resp.) ferrets, each weighing between 1.5 - 2.1 kg.
- Surgery and Animal Preparation: The techniques involved in the surgery and preparation for recording are described in detail in Shamma et al. (1993). The ferrets were anesthetized with pentobarbital sodium (40 mg/kg IP) and maintained in an areflexic state using a continuous IV infusion of pentobarbital (~ 5 mg/kg/hr) diluted with dextrose-electrolyte solution for metabolic stability. Data collection typically lasted 60-70 hours.
- Recording Procedures: Single-unit action potentials were recorded using glass-insulated tungsten microelectrodes with 5 to 6 M Ω impedances. The recorded signals were led through amplifiers and filters. Depending on the paradigm, a stimulus was presented every few seconds, and raster plots with 1ms time resolution were produced.
- In AI, recordings were typically made at depths of 300 - 600 μ m (layers III and IV). In AAF, electrode penetrations were made parallel to the depth of the suprasylvian sulcus (SSS), approximately 0.5 mm caudal of the sulcus so that cortical layers III and/or IV were reached. IC was exposed by removal of (visual) cortex, and electrodes were lowered until ICC was reached, following standard criteria.


Change in Ripple Phase




Measuring the Ripple Transfer Function



← Tonotopic Axis →



Amplitude





Phase

Ripple Frequency (cyc/oct)


0.8 cycles/octave
at 8 different phases

Stimulus onset: 100 ms
Stimulus offset: 150 ms


0 phase ripple at
6 different ripple freq.




Tuning as a function of Ripple Frequency and Phase




Response fields of varying bandwidth




Response Fields with varying asymmetries




The Anterior Cortical Field




CAT







FERRET







Area of
magnification




BF's and latencies are similar



Bandwidths (BW20) are larger in AAF




Tuning to Ripples tends to be lower in AAF





Ω_0 = Best
Stationary
Ripple
Frequency
(cyc/oct)

\square Single Unit
 \blacksquare Cluster




RF's asymmetries are evenly distributed




Predictions Using Stationary Ripples


**Response Field
of Cell**




**Spectral Profile
of stimulus**




**Response of Cell
to Profile**







The Inferior Colliculus







R
L






Of BF's and Latencies



Bandwidths





Best Ripple Frequency





Ω_0 = Best Ripple Frequency

RF's are more symmetrical in IC



□ Single Unit
▨ Cluster

Temporal properties: Moving Ripples




- Ω = ripple frequency in cycle/octaves
- ω = temporal frequency in Hz

Responses to Moving Ripples

Ripple Frequency is 0.4 cycles/oct

Temporal Frequency (Hz)




$\Omega = 0.4$ cyc/oct

$\omega = 4$ to 32 Hz



Time (ms)

30 sweeps per ω

Step 1. From Spike Count to Period Histogram



Step 2. Magnitude and Phase of Best Fit




Impulse Response Function

Linearity of responses in the Auditory Cortex

AI and AAF display similar characteristics


Anterior Field:

- (A) Stimulus Spectrogram, $S(x,t)$ (B) $RF(x)$ (C) $\sum_x RF(x) \cdot S(x,t)$ (D) $IR(t)$ (E) Predicted Response





Inferior Colliculus Responses



8
24
40
56

Time (ms)

$\Omega = 0.2$ cyc/oct

15 sweeps per ω

$\omega = 4$ to 64 Hz

Summary

- Stationary and traveling ripples can be used to extract spectral and temporal properties of auditory cortical neurons.
- Linearity: Responses to a broad-band complex stimuli, decomposed into a linear combination of ripples, can be predicted by summing the neuronal responses to the individual ripples.
- Only Cortical neurons are selective to ripple frequencies; Collicular neurons are low-pass with respect to ripple frequencies.
- Therefore, AI and AAF neurons could perform a multi-scale analysis of spectral shape: the spectral profile is analyzed at different degrees of resolution by neurons with receptive fields of different best frequencies, bandwidths and asymmetries.

References

- DeValois R. and DeValois K. (1988) *Spatial Vision*. New York: Oxford U. Press
- Shamma S.A., Versnel H. and Kowalski N. (1995) "Ripple analysis in ferret primary auditory cortex I. Response characteristics of single units to sinusoidally rippled spectra". *Auditory Neuroscience* 1(3), pp. 233-254.
- Shamma S.A. and Versnel H. (1995) "Ripple analysis in ferret primary auditory cortex. II. Prediction of unit responses to arbitrary spectral profiles". *Auditory Neuroscience* 1(3), pp. 255-270.
- Versnel H., Kowalski N. and Shamma S.A. (1995) "Ripple analysis in ferret primary auditory cortex.III. Topographic distribution of ripple response parameters". *Auditory Neuroscience* 1(3), pp. 271-285.
- Schreiner C.E. and Calhoun B.M. (1995) "Spectral envelope coding in cat primary auditory cortex: properties of ripple transfer functions". *Auditory Neuroscience* 1(1): 23 pages.
- Kowalski N., Versnel H. and Shamma S.A. (1995) "Comparison of responses in the anterior and primary auditory fields of the ferret cortex". *J. Neurophys.* 73(4), pp.1513-1523.
- Owens A. and Shamma S. "Surface evoked potentials reveal selectivity to spectral shape features". ARO 96.
- Kowalski N., D.A.D. and Shamma S.A. (1995) "Analysis of dynamic spectra in ferret primary auditory cortex: I. Response characteristics of single units to moving rippled spectra".
- Kowalski N., D.A.D. and Shamma S.A. (1995) "Analysis of dynamic spectra in ferret primary auditory cortex: II. Prediction of unit responses to arbitrary dynamic spectra".